

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7283: Recombinant Human Carbonic Anhydrase 10/CA10 (C-6His)

Gene ID: CA10
Gene ID: 56934
Uniprot ID: Q9NS85

## **Description**

Source: Human Cells. MW :32.82kD.

Recombinant Human Carbonic Anhydrase 10 is produced by our Mammalian expression system and the target gene encoding Gln22-Asn300 is expressed with a 6His tag at the C-terminus. Carbonic Anhydrase X (CA10) belongs to CA family of zinc metalloenzymes, which catalyze the reversible hydration of carbon dioxide in various biological processes such as respiration, renal tubular acidification and bone resorption. While CA10 is a secreted protein without Carbonic Anhydrase activity (i.e., the reversible hydration of CO2) due to point mutations in the zinc binding site, it has esterase activity. The human and mouse CA10 are expressed in the brain, indicating that they may play a role in brain development.

## **Product Info**

**Amount :** 10 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM TrisHCl, 150mM NaCl, pH 8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: QQNSPKIHEGWWAYKEVVQGSFVPVPSFWGLVNSAWNLCSVGKRQSPVNIETSHMIFDPFLTPLRINTGGRKV

 $SGTMYNTGRHVSLRLDKEHLVNISGGPMTYSHRLEEIRLHFGSEDSQGSEHLLNGQAFSGEVQLIHYNHELYTN\\ VTEAAKSPNGLVVVSIFIKVSDSSNPFLNRMLNRDTITRITYKNDAYLLQGLNIEELYPETSSFITYDGSMTIPPCYE$ 

TASWIIMNKPVYITRMQMHSLRLLSQNQPSQIFLSMSDNFRPVQPLNNRCIRTNVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.