

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7318: Recombinant Human Insulin-Like Growth Factor-Binding Protein 4/IGFBP-4 (C-6His)

Gene ID: 3487
Uniprot ID: P22692

## **Description**

Source: Human Cells. MW ·27 01kD

Recombinant Human Insulin-Like Growth Factor-Binding Protein 4 is produced by our Mammalian expression system and the target gene encoding Asp22-Glu258 is expressed with a 6His tag at the C-terminus. IGFBP-4, whose full name is insulin-like growth factor-binding protein 4, is induced by forskolin and N6, O2'dibutyryl sdenosine 3', or 5'-cyclic monophosphate. It contains IGFBP N-terminal domain and thyroglobulin type-1 domain, and can bind IGF2. The IGF-binding proteins can prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture.

## **Product Info**

Amount:  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: DEAIHCPPCSEEKLARCRPPVGCEELVREPGCGCCATCALGLGMPCGVYTPRCGSGLRCYPPRG

VEKPLHTLMHGQGVCMELAEIEAIQESLQPSDKDEGDHPNNSFSPCSAHDRRCLQKHFAKIRDRS TSGGKMKVNGAPREDARPVPQGSCQSELHRALERLAASQSRTHEDLYIIPIPNCDRNGNFHPKQC

HPALDGQRGKCWCVDRKTGVKLPGGLEPKGELDCHQLADSFREVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\hat{A}\mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.