

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7328: Recombinant Human Junctional Adhesion Molecule B/JAM-B/CD322 (C-6His)

**Gene ID**: 58494 **Uniprot ID**: P57087

## **Description**

Source: Human Cells. MW ·24 29kD

Recombinant Human JAM-B is produced by our Mammalian expression system and the target gene encoding Phe29-Asn236 is expressed with a 6His tag at the C-terminus. Junctional Adhesion Molecule B (JAM-B) is a single-pass type I membrane protein that belongs to the juctional adhesion molecules family. JAM-B includes a signal sequence (aa 1-28), an extracellular region (aa 29-238) with one Ig-like C2-type domain and one Ig-like V-type domain, a transmembrane segment (aa 239-259), and a cytoplasmic domain (aa 260 - 298). JAMB is localized to the tight junctions between endothelial cells or epithelial cells. JAM-B is prominently expressed in the heart, placenta, lung, foreskin and lymph node. It is also present on the endothelia of other vessels. JAM-B acts as an adhesive ligand for interacting with a variety of immune cell types and may play a role in lymphocyte homing to secondary lymphoid organs.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition:** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: FSAPKDQQVVTAVEYQEAILACKTPKKTVSSRLEWKKLGRSVSFVYYQQTLQGDFKNRAEMIDFNI

RIKNVTRSDAGKYRCEVSAPSEQGQNLEEDTVTLEVLVAPAVPSCEVPSSALSGTVVELRCQDKE GNPAPEYTWFKDGIRLLENPRLGSQSTNSSYTMNTKTGTLQFNTVSKLDTGEYSCEARNSVGYRR

**CPGKRMQVDDLNVDHHHHHH** 

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\hat{A}\mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.