

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7361: Recombinant Human Wnt Inhibitory Factor 1/WIF-1 (C-6His)

Gene : WIF1 **Gene ID :** 11197 **Uniprot ID :** 09Y5W5

Description

Source: Human Cells. MW:39.47kD.

Recombinant Human Wnt Inhibitory Factor 1 is produced by our Mammalian expression system and the target gene encoding Gly29-Trp379 is expressed with a 6His tag at the C-terminus. Wnt Inhibitory Factor 1 (WIF1) is a secreted protein, which binds WNT proteins and inhibits their activities. WNT proteins are extracellular signaling molecules involved in the control of embryonic development. WIF1 contains a WNT inhibitory factor (WIF) domain and 5 epidermal growth factor (EGF)-like domains. is found to be present in fish, amphibia and mammals. WIF1 is a recurrent target in human salivary gland oncogenesis.WIF1 may be involved in mesoderm segmentation. WIF1 is a tumor suppressor, specifically in nonfunctioning pituitary tumors.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 μm filtered solution of 10mM HAc-NaAc, 150mM NaCl, 0.5% CHAPS, pH

4.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: GPPQEESLYLWIDAHQARVLIGFEEDILIVSEGKMAPFTHDFRKAQQRMPAIPVNIHSMNFTWQAAGQAEYFYE

FLSLRSLDKGIMADPTVNVPLLGTVPHKASVVQVGFPCLGKQDGVAAFEVDVIVMNSEGNTILKTPQNAIFFKTC QQAECPGGCRNGGFCNERRICECPDGFHGPHCEKALCTPRCMNGGLCVTPGFCICPPGFYGVNCDKANCSTT CFNGGTCFYPGKCICPPGLEGEQCEISKCPQPCRNGGKCIGKSKCKCSKGYQGDLCSKPVCEPGCGAHGTCHE

PNKCQCQEGWHGRHCNKRYEASLIHALRPAGAQLRQHTPSLKKAEERRDPPESNYIWVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \Box \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.