

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7371: Recombinant Human Dickkopf-Related Protein 3/DKK3 (C-6His)(Discontinued)

Gene ID: 27122 Uniprot ID: Q9UBP4

Description

Source: Human Cells. MW:37.22kD.

Recombinant Human Dickkopf-Related Protein 3 is produced by our Mammalian expression system and the target gene encoding Ala22-Ile350 is expressed with a 6His tag at the C-terminus. Dickkopf-related protein 3 (DKK3) belongs to the DKK protein family including Dkk-1, 2, 3 and -4. DKK3 is a 350 amino acid secreted glycoprotein which is comprised of an N-terminal signal peptide and 2 conserved cysteine-rich domains that are separated by a 12 amino acid linker region. Dkk-3 also have one prokineticin domain. DKK3 is involved in embryonic development through its inhibition of the WNT signaling pathway. The Dkk family also includes Soggy, which is homologous to Dkk-3 but not to the other family members. Soggy has not been shown to inhibit Wnt signaling, and its role in the pathway is unclear.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: APAPTATSAPVKPGPALSYPQEEATLNEMFREVEELMEDTQHKLRSAVEEMEAEEAAAKASSEVNLANLPPSYH

NETNTDTKVGNNTIHVHREIHKITNNQTGQMVFSETVITSVGDEEGRRSHECIIDEDCGPSMYCQFASFQYTCQ PCRGQRMLCTRDSECCGDQLCVWGHCTKMATRGSNGTICDNQRDCQPGLCCAFQRGLLFPVCTPLPVEGEL CHDPASRLLDLITWELEPDGALDRCPCASGLLCQPHSHSLVYVCKPTFVGSRDQDGEILLPREVPDEYEVGSFM

EEVROELEDLERSLTEEMALGEPAAAAAALLGGEEIVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.