

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7386: Recombinant Human Apolipoprotein H/ApoH (C-6His)

**Gene ID:** 350 **Uniprot ID:** P02749

## **Description**

Source: Human Cells. MW:37.29kD.

Recombinant Human Apolipoprotein H is produced by our Mammalian expression system and the target gene encoding Gly20-Ser345 is expressed with a 6His tag at the C-terminus. Apolipoprotein H (ApoH) is a 50 kDa variably glycosylated member of the complement control superfamily of proteins. Human ApoH is a major phospholipid binding protein and an important component to measure in the assessment of anti-phospholipid syndrome. Hepatocyte-derived ApoH binds to negatively charged phospholipids. It circulates as a component of lipoprotein particles and as a lipid-free serum protein. Human ApoH is also more specific than anti-cardiolipin antibodies and its presence correlates better with thrombotic risk. Mature human ApoH shares 76% and 82% aa sequence identity with mouse and rat ApoH.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: GRTCPKPDDLPFSTVVPLKTFYEPGEEITYSCKPGYVSRGGMRKFICPLTGLWPINTLKCTPRVCPFAGILENGAV

RYTTFEYPNTISFSCNTGFYLNGADSAKCTEEGKWSPELPVCAPIICPPPSIPTFATLRVYKPSAGNNSLYRDTAVF ECLPQHAMFGNDTITCTTHGNWTKLPECREVKCPFPSRPDNGFVNYPAKPTLYYKDKATFGCHDGYSLDGPEEI ECTKLGNWSAMPSCKASCKVPVKKATVVYQGERVKIQEKFKNGMLHGDKVSFFCKNKEKKCSYTEDAQCIDG

TIEVPKCFKEHSSLAFWKTDASDVKPCVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.