

32-7387: Recombinant Human EMMPRIN/Basigin/CD147 (C-6His)

Gene : BSG
Gene ID : 682
Uniprot ID : P35613

Description

Source: Human Cells.

MW :21.16kD.

Recombinant Human EMMPRIN is produced by our Mammalian expression system and the target gene encoding Ala22-His205 is expressed with a 6His tag at the C-terminus. Extracellular Matrix Metalloproteinase Inducer (EMMPRIN) belongs to the immunoglobulin superfamily, which has the homology to both the immunoglobulin V domain and MHC class II antigen beta-chain. EMMPRIN is a transmembrane glycoprotein with different forms, resulting from different modes of glycosylation and N-terminal sequence variants. EMMPRIN can be expressed in breast cancer, oral squamous cell carcinoma, glioma, lymphoma, lung, bladder, and melanoma carcinomas cells. EMMPRIN promotes invasion, metastasis, growth, and survival of malignants cells, serves as a receptor for extracellular cyclophilinthe, may play a role in signal transduction.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : AAGTVFTTVEDLGSKILLTCSLNDSEVTGHRWLKGGVVLKEDALPGQKTEFKVDSDDQWGEYSCVFLPEPM
GTANIQLHGPPRVKAVKSSEHINEGETAMLVCKSESVPVTDWAWYKITDSEDKALMNGSESRRFFVSSSQGRS
ELHIENLNMEADPGQYRCNGTSSKGSDQAIITLRVRSHVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.