

## 32-7399: Recombinant Human Sialic Acid Binding Ig-Like Lectin 3/Siglec-3/CD33 (C-Fc-6His)

**Gene :** CD33  
**Gene ID :** 945  
**Uniprot ID :** P20138

### Description

Source: Human Cells.  
MW :55.01kD.

Recombinant Human Sialic Acid Binding Ig-like Lectin 3 is produced by our Mammalian expression system and the target gene encoding Asp18-His259 is expressed with a Fc, 6His tag at the C-terminus. CD33 is a type I Lectin belonging to the Ig superfamily. CD33 contains an N terminal Ig like V type domain, which mediates sialic acid binding, followed by one Ig like C2 type domain, a transmembrane region and a cytoplasmic tail containing two conserved immunoreceptor tyrosine based inhibition motifs (ITIMs). Eleven human Siglecs have been characterized. Siglecs 5 to 11 share a high degree of sequence similarity with CD33/Siglec3 both in their extracellular and intracellular regions. They are collectively referred to as CD33 related Siglecs. CD33 related Siglecs have differential expression pattern within the hematopoietic system. They are involved in the regulation of cellular activation within the immune system. Siglec 3 expression is restricted to cells of myelomonocytic lineage. Siglec3 recruits SHP1 and SHP2 to its ITIMs upon phosphorylation.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 2mM EDTA, pH 7.2.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** DPNFWLQVQESVTVQEGLCVLPCTFFHPIPYDKNSPVHGYWFREGAIISGDSPVATNKLDQEVQEETQGRF  
RLLGDPSRNNCSLSIVDARRRDNGSYFFRMERGSTKYSYKSPQLSVHVTDLTHRPKILIPGTLEPGHSKNLTCSV  
SWACEQGTPIFSWLSAAPTSLGPRTHSSVLIITPRPQDHGTNLTCQVKFAGAGVTTERTIQLNVTVYPQNPTT  
GIFPGDGSGKQETRAGVVHTSDIEGRMDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCV  
VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKT  
ISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKL  
TVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKVDHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.