

32-7408: Recombinant Human Clusterin/ApoJ (C-6His)

Gene : CLU
Gene ID : 1191
Uniprot ID : P10909

Description

Source: Human Cells.
MW :51.1kD.

Recombinant Human Clusterin is produced by our Mammalian expression system and the target gene encoding Asp23-Glu449 is expressed with a 6His tag at the C-terminus. Clusterin is a secreted protein which belongs to the Clusterin family. Clusterin is expressed in adult testis, heart, ovary, adrenal gland, brain and liver. Clusterin has been suggested to be involved in several basic biological events such as cell death, tumor progression, and neurodegenerative disorders. In addition, Clusterin is up/ down regulated on the mRNA or protein level in many pathological and clinically relevant situations including cancer, organ regeneration, infection, Alzheimer disease, retinitis pigmentosa, myocardial infarction, renal tubular damage, autoimmunity and others.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : DQTVSDNELQEMSNQGSKYVNKEIQNAVNGVKQIKTLIEKTNEERKTLLSNLEEAKKKKEDALNETRESETKLKE
LPGVCNETMMALWEECKPCLKQTCMKFYARVCRSGSLVGRQLEEFNLQSSPFYFWMNGDRIDSLLENDRQ
QTHMLDVMQDHFSRASSIIDELFQDRFFTRPQDQTYHYLPFSLPHRRPHFFPKSRIVRSLMPFSPYEPLNPFHAM
FQPFLEMIHEAQQAMDIHFHSPAQHPPTFIREGDDDRVTVCREIRHNSTGCLRMKDQCDKCREILSVDCSTNN
PSQAKLRRELDSESLQVAERLTRKYNELLKSYQWKMLNTSSLEQLNEQFNWVSRLANLTQGEDQYYLRVTTVA
SHTSDSDVPSGVTEVVVKLFDSDPITVTPVEVSRKNPKFMETVAEKALQEYRKKHREEVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.