

32-7413: Recombinant Human Tissue Inhibitor of Metalloproteinases 2/TIMP-2 (C-6His)

 Gene :
 TIMP2

 Gene ID :
 7077

 Uniprot ID :
 P16035

Description

Source: Human Cells.

MW :22.79kD.

Recombinant Human Tissue Inhibitor of Metalloproteinases 2 is produced by our Mammalian expression system and the target gene encoding Cys27-Pro220 is expressed with a 6His tag at the C-terminus. Tissue Inhibitor of Metalloproteinases 2 (TIMP-2) complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them by binding to their catalytic zinc cofactor. Known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-10, MMP-13, MMP-14, MMP-15, MMP-16 and MMP-19.

Product Info

Amount :	10 μg / 50 μg
Content :	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition :	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid :	CSCSPVHPQQAFCNADVVIRAKAVSEKEVDSGNDIYGNPIKRIQYEIKQIKMFKGPEKDIEFIYTAPSSAVCGVSL DVGGKKEYLIAGKAEGDGKMHITLCDFIVPWDTLSTTQKKSLNHRYQMGCECKITRCPMIPCYISSPDECLWMD WVTEKNINGHQAKFFACIKRSDGSCAWYRGAAPPKQEFLDIEDPVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 \tilde{A} $\hat{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ \tilde{A} \square $\hat{A}\mu$ g (1 IEU/ \tilde{A} \square $\hat{A}\mu$ g) as determined by LAL test.