

32-7417: Recombinant Human Preadipocyte Factor-1/Protein d Homolog 1/Pref-1/DLK-1 (C-6His)

Gene : DLK1
Gene ID : 8788
Uniprot ID : P80370

Description

Source: Human Cells.

MW :30.15kD.

Recombinant Human DLK-1 is produced by our Mammalian expression system and the target gene encoding Ala24-Pro297 is expressed with a 6His tag at the C-terminus. Protein Delta Homolog 1 (DLK-1) is a transmembrane protein which contains a signal peptide, an extracellular domain with six tandem epidermal growth factor (EGF)-like domains, a single pass transmembrane domain, and a short cytoplasmic tail. It is found within the stromal cells in close contact to the vascular structure of placental villi, yolk sac, fetal liver, adrenal cortex and pancreas and in the beta cells of the islets of Langerhans in the adult pancreas. In addition, it is detected in some forms of neuroendocrine lung tumor tissue. DLK-1 may have a important role in neuroendocrine differentiation.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : AECFPACNPQNGFCEDDNVCRCQPGWQGPLCDQCVTSPGCLHGLCGEPGQCICITDGWDGELCDRDVRACS
SAPCANNGTCVSLDDGLYECSCAPGYSGKDCQKKDGPVINGSPCQHGGTCVDDEGRASHASCLCPPGFSG
NFCEIVANSCTPNPCENDGVCTDIGGDFRCRCPAGFIDKTCRPTVNCASSPCQNGGTCLQHTQVSYECLCKP
EFTGLTCVKKRALSPQQVTRLPNGYGLAYRLTPGVHELPVQQPEHRILKVSMEKLNKTPVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.