

## 32-7428: Recombinant Human Glycoprotein A33/GPA33/GPA33 (C-6His)

**Gene :** GPA33

**Gene ID :** 10223

**Uniprot ID :** Q99795

### Description

Source: Human Cells.

MW :24.66kD.

Recombinant Human Glycoprotein A33 is produced by our Mammalian expression system and the target gene encoding Ile22-Val235 is expressed with a 6His tag at the C-terminus. Human Glycoprotein A33 (GPA33) is a single-pass type I membrane protein, belongs to the CTX family of cell adhesion molecular within the immunoglobulin family, can be expressed in normal gastrointestinal epithelium and in 95% of colon cancers. GPA33 consists of one Ig-like C2-type domain and one Ig-like V-type domain. The predicted mature protein includes a single transmembrane domain, a extracellular region and a intracellular tail. Intracellular traffic and recycling to the cell surface appear to play an important role in GPA33 function and to have an influence on its surface density superseding translation regulation. GPA33 has become a promising target of immunologic therapy strategies. GPA33 may also play a important role in cell-cell recognition and signaling.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** ISVETPQDVLRASQGKSVTLPCYHTSTSSREGLIQWDKLLLTHTERVVIWPFSSNKNYIHGELYKNR  
VSISNNAEQSDASITIDQLTMADNGTYECVSLMSDLEGNTKSRVRLLVLPVPPSKPECGIEGETIIGN  
NIQLTCQSKEGSPTQYSWKRYNILNQEQLAQPASGQPVSLKNISTDTSGYYICTSSNEEGTQFC  
NITVAVRSPSMNVVDHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.