

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7434: Recombinant Human Ephrin-A4/EFNA4 (C-Fc-6His)

Gene ID: 1945 Uniprot ID: P52798

Description

Source: Human Cells.

MW:44.3kD.

Recombinant Human Ephrin-A4 is produced by our Mammalian expression system and the target gene encoding Leu26-Gly171 is expressed with a Fc, 6His tag at the C-terminus. Ephrin-A4 is a member of the ephrin ligand family which binds members of the Eph receptor family. All ligands share a conserved extracellular sequence, which most likely corresponds to the receptor binding domain. Ephrin-A4 consists of approximately 125 amino acids and includes four invariant cysteines, It has been shown to bind EphA2, EphA3, EphA4, EphA5, EphA6, EphA7, and EphB1. Ephrin-A4 binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. It may play a role in the interaction between activated B-lymphocytes and dendritic cells in tonsils.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: LRHVVYWNSSNPRLLRGDAVVELGLNDYLDIVCPHYEGPGPPEGPETFALYMVDWPGYESCQAEGPRAYKRW

VCSLPFGHVQFSEKIQRFTPFSLGFEFLPGETYYYISVPTPESSGQCLRLQVSVCCKERKSESAHPVGSPGESGV DDIEGRMDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSR EEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSV

MHEALHNHYTQKSLSLSPGKHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than $0.1 \text{ ng}/\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.