

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7459: Recombinant Human Neurotrophic Tyrosine Kinase Receptor Type 2/TrkB/NTRK2 (C-6His)

Gene ID: NTRK2
Gene ID: 4915
Uniprot ID: Q16620

Description

Source: Human Cells. MW :45.25kD.

Recombinant Human TrkB is produced by our Mammalian expression system and the target gene encoding Cys32-His430 is expressed with a 6His tag at the C-terminus. The TRK Family of Tyrosine Kinase Receptor consists of 3 members: TrkA, TrkB and TrkC. The three TRK family proteins have different ligand specificities. They connect to different neurotrophins, including NGF, BDNF, NT-3NT-4/5. TRKA binds NGF, TRKB binds BDNF and NT-3, TRKC binds NT-4/5. At the protein sequence level, human and rat TRKB have greater than 90% sequence identity and the proteins exbihit cross-species activity. TRKB is primarily expressed in the nervous system and it also expression in a wide variety of tissues with low levels.

Product Info

Amount: 10 μg / 50 μg

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20 $^{\circ}\text{C},$ though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: CPTSCKCSASRIWCSDPSPGIVAFPRLEPNSVDPENITEIFIANQKRLEIINEDDVEAYVGLRNLTIVDSGLKFVAH

KAFLKNSNLQHINFTRNKLTSLSRKHFRHLDLSELILVGNPFTCSCDIMWIKTLQEAKSSPDTQDLYCLNESSKNI PLANLQIPNCGLPSANLAAPNLTVEEGKSITLSCSVAGDPVPNMYWDVGNLVSKHMNETSHTQGSLRITNISSD DSGKQISCVAENLVGEDQDSVNLTVHFAPTITFLESPTSDHHWCIPFTVKGNPKPALQWFYNGAILNESKYICTK IHVTNHTEYHGCLQLDNPTHMNNGDYTLIAKNEYGKDEKQISAHFMGWPGIDDGANPNYPDVIYEDYGTAAND

IGDTTNRSNEIPSTDVTDKTGREHVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.