

32-7522: Recombinant Human CD8 beta Chain/CD8B (C-6His)

Gene : CD8B
Gene ID : 926
Uniprot ID : P10966

Description

Source: Human Cells.

MW :17.8kD.

Recombinant Human CD8B is produced by our Mammalian expression system and the target gene encoding Leu22-Pro170 is expressed with a 6His tag at the C-terminus. T-Cell Surface Glycoprotein CD8 beta Chain (CD8 Antigen) is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. CD8 Antigen, acting as a coreceptor, and the T-cell receptor on the T lymphocyte recognize antigens displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. Multiple alternatively spliced transcript variants encoding distinct membrane associated or secreted isoforms have been described. A pseudogene, also located on chromosome 2, has been identified.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : LQQTPAYIKVQTNKMVMSCEAKISLSNMRIYWLQRQAPSSDSHHEFLALWDSAKGTIHGEEVEQEKIIVFRD
ASRFILNLTSVKPEDSGIYFCMIVGSPELTFGKGTQLSVVDLPTTAQPTKKSTLKKRVCRLPRPETQKGPLCSPV
DHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.