

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7545: Recombinant Human Leucine-Rich Repeat-Containing Protein 2/LRRN2 (C-6His)(Discontinued)

Gene ID: LRRTM2
Gene ID: 26045
Uniprot ID: O43300

Description

Source: Human Cells. MW:45.55kD.

Recombinant Human LRRTM2 is produced by our Mammalian expression system and the target gene encoding Cys34-Arg422 is expressed with a 6His tag at the C-terminus. Leucine-Rich Repeat Transmembrane Neuronal Protein 2 (LRRTM2) is a single-pass type I membrane protein that belongs to the LRRTM family. It contains ten LRR (leucine-rich) repeats, one LRRCT domain, and one LRRNT domain. LRRTM2 is expressed in neuronal tissues, and it interacts with DLG4 and NRXN1. LRRTM2 has been suggested to be involved in the development and maintenance of excitatory synapses in the vertebrate nervous system. LRRTM2 also regulates the surface expression of AMPA receptors. LRRTM2 acts as a ligand for the presynaptic receptors NRXN1-A and NRXN1-B.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: CPPKCRCEKLLFYCDSQGFHSVPNATDKGSLGLSLRHNHITELERDQFASFSQLTWLHLDHNQIST

VKEDAFQGLYKLKELILSSNKIFYLPNTTFTQLINLQNLDLSFNQLSSLHPELFYGLRKLQTLHLRSNS LRTIPVRLFWDCRSLEFLDLSTNRLRSLARNGFAGLIKLRELHLEHNQLTKINFAHFLRLSSLHTLFLQ WNKISNLTCGMEWTWGTLEKLDLTGNEIKAIDLTVFETMPNLKILLMDNNKLNSLDSKILNSLRSLTT VGLSGNLWECSARICALASWLGSFQGRWEHSILCHSPDHTQGEDILDAVHGFQLCWNLSTTVTVM

ATTYRDPTTEYTKRISSSSYHVGDKEIPTTAGIAVTTEEHFPEPDNAIFTQRVDHHHHHH

Application Note

Storage condition:

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\hat{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.