

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7568: Recombinant Mouse Trefoil Factor 3/TFF3 (C-6His)

Gene : Tff3 **Gene ID :** 21786 **Uniprot ID :** Q62395

Description

Source: Human Cells.

MW:7.3kD.

Recombinant Mouse Trefoil Factor 3 is produced by our Mammalian expression system and the target gene encoding Ala23-Phe81 is expressed with a 6His at the C-terminus. Trefoil factors (TFF) are secretory products of mucin producing cells. They play a key role in the maintenance of the surface integrity of oral mucosa and enhance healing of the gastrointestinal mucosa by a process called restitution. TFF comprises the gastric peptides (TFF1), spasmolytic peptide (TFF2), and the intestinal trefoil factor (TFF3). They have an important and necessary role in epithelial restitution within the gastrointestinal tract. Members of the trefoil family are characterized by having at least one copy of the trefoil motif, a 40-amino acid domain that contains three conserved disulfide bonds. They are stable secretory proteins expressed in gastrointestinal mucosa. Trefoil Factor 3(TFF3) is involved in the maintenance and repair of the intestinal mucosa. TFF3 promotes the mobility of epithelial cells in healing processes (motogen).

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: ADYVGLSPSQCMVPANVRVDCGYPSVTSEQCNNRGCCFDSSIPNVPWCFKPLQETECTFHHHHH

Н

Application Note

Storage condition:

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\hat{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.