9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-7580: Recombinant Human Integral Membrane Protein 2B/ITM2B/imBRI2 (C-6His)

Gene : ITM2B
Gene ID : 9445
Uniprot ID : Q9Y287

Description

Source: Human Cells.
MW :23.3kD.
Recombinant Human Integral Membrane Protein 2B is produced by our Mammalian expression system and the target gene encoding Tyr76-Ser266 is expressed with a 6His tag at the C-terminus. Integral Membrane Protein 2B (ITM2B) is expressed in the Golgi and on the cell surface. ITM2B forms homodimer through disulfide-linked interaction with SPPL2A, SPPL2B and APP. ITM2B is expressed in brain and the other tissues. Defects in ITM2B cause cerebral amyloid angiopathy ITM2B-related type 1(CAA-ITM2B1) and amyloid angiopathy ITM2B-related type 2(CAA-ITM2B2). CAA-ITM2B1 is characterized by amyloid deposition in the walls of cerebral blood vessels and neurodegeneration in the central nervous system. CAA-ITM2B2 characterized by amyloid deposition in the walls of the blood vessels of the cerebrum, choroid plexus, cerebellum, spinal cord and retina.

Product Info

Amount :	10 µg / 50 µg
Content :	Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition :	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid :	YKYFALQPDDVYYCGIKYIKDDVILNEPSADAPAALYQTIEENIKIFEEEEVEFISVPVPEFADSDPANIVHDFNKKL TAYLDLNLDKCYVIPLNTSIVMPPRNLLELLINIKAGTYLPQSYLIHEHMVITDRIENIDHLGFFIYRLCHDKETYKLQ RRETIKGIQKREASNCFAIRHFENKFAVETLICSVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 ÃÂµg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Endotoxin : Less than 0.1 ng/ÃÂµg (1 IEU/ÃÂµg) as determined by LAL test.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.