

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7585: Recombinant Human Leucine-Rich Repeat-Containing Protein 3B/LRRC3B (C-6His)

Gene ID: LRRC3B
Gene ID: 116135
Uniprot ID: Q96PB8

## **Description**

Source: Human Cells.

MW:20kD.

Recombinant Human LRRC3B is produced by our Mammalian expression system and the target gene encoding Cys34-Tyr204 is expressed with a 6His tag at the C-terminus. Leucine-Rich Repeat-Containing Protein 3B (LRRC3B) belongs to the LRRC3 family. LRRC3B is single-pass membrane protein and contains three leucine-rich repeats, one LRRCT domain, and one LRRNT domain. LRR-containing proteins, of which there are greater than 2,000, participate in many important processes, including plant and animal immunity, hormone-receptor interactions, cell adhesion, signal transduction, regulation of gene expression, and apoptosis. A number of microarray expression profiling studies on human cancers have shown that LRRC3B is down-regulated in gastric, breast, colon, testis, prostate and brain cancers, suggesting LRRC3B involvement in carcinogenesis

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM Tris,150mM NaCl,pH8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition:** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: CPKGCLCSSSGGLNVTCSNANLKEIPRDLPPETVLLYLDSNQITSIPNEIFKDLHQLRVLNLSKNGIEFIDEHAFKG

VAETLQTLDLSDNRIQSVHKNAFNNLKARARIANNPWHCDCTLQQVLRSMASNHETAHNVICKTSVLDEHAGR

PFLNAANDADLCNLPKKTTDYVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\tilde{A} \square \hat{A} \mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$  (1 IEU/ $\tilde{A} \square \hat{A} \mu g$ ) as determined by LAL test.