

32-7590: Recombinant Human IL-6 Receptor Subunit α /IL-6RA/CD126 (C-6His)

Gene : IL6R
Gene ID : 3570
Uniprot ID : P08887

Description

Source: Human Cells.
MW :38.9kD.

Recombinant Human Interleukin-6 receptor alpha is produced by our Mammalian expression system and the target gene encoding Leu20-Asp358 is expressed with a 6His tag at the C-terminus. Interleukin 6 is a potent pleiotropic cytokine that regulates cell growth and differentiation and plays an important role in the immune response. IL6Ra is a part of the receptor for interleukin 6 cytokine. IL6Ra binds to IL6 with low affinity, but does not transduce a signal. Signal activation necessitates an association with IL6ST. Activation may lead to the regulation of the immune response, acute-phase reactions and hematopoiesis. Low concentration of a soluble form of IL6 receptor acts as an agonist of IL6 activity. Dysregulated production of IL6 and this receptor are implicated in the pathogenesis of many diseases, such as multiple myeloma, autoimmune diseases and prostate cancer.

Product Info

Amount : 10 μ g / 50 μ g
Content : Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : LAPRRCPAQEVARGVLTSLPGDSVLTCTPGVEPEDNATVHWVLRKPAAGSHPSRWAGMGRRLLLRSVQLHDS
GNYSCYRAGRPAQTVHLLVDVPPEEPQLSCFRKSPLSNVCEWGRSTPSLTTKAVLLVRKFQNSPAEDFQEP
QYSQESQKFSCQLAVPEGDSSFYIVSMCVASSVGSKFSTQTFQGCILQPDPPANITVTAVARNPRWLSVTW
QDPHSWNSSFYRLRFELRYAERSKTFTTWVMDLQHHCVIHDWGLRHVVQLRAQEEFGQGEWSEWSPE
AMGTPWTESRSPPAENEVSTPMQALTTNKDDDNILFRDSANATSLPVQDQDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ μ g (1 IEU/ μ g) as determined by LAL test.