

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7593: Recombinant Mouse Tumor Necrosis Factor Receptor II/TNFRSF1B/CD120b (C-6His)

Gene ID: 21938
Uniprot ID: Q545P4

## **Description**

Source: Human Cells.

MW:26.4kD.

Recombinant Mouse Tumor Necrosis Factor Receptor II is produced by our Mammalian expression system and the target gene encoding Val23-Gly258 is expressed with a 6His tag at the C-terminus. Tumor Necrosis Factor Receptor Superfamily Member 1B (TNFRSF1B) is a member of the Tumor Necrosis Factor Receptor Superfamily. TNFRSF1B contains four TNFR-Cys repeats. TNFRSF1B can be cleaved into the following 2 chains: Tumor necrosis factor receptor superfamily member 1b and membrane form and Tumor necrosis factor-binding protein 2. TNFRSF1B is a receptor with high affinity for TNFSF2/TNF-a and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-a. TNFRSF1B mediates most of the metabolic effects of TNF-a. TNF-a-induced apoptosis suggests that it regulates TNF-a function by antagonizing its biological activity.

## **Product Info**

**Amount:**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: VPAQVVLTPYKPEPGYECQISQEYYDRKAQMCCAKCPPGQYVKHFCNKTSDTVCADCEASMYTQ

VWNQFRTCLSCSSSCTTDQVEIRACTKQQNRVCACEAGRYCALKTHSGSCRQCMRLSKCGPGFGVASSRAPNGNVLCKACAPGTFSDTTSSTDVCRPHRICSILAIPGNASTDAVCAPESPTLSAIPRTL

YVSQPEPTRSQPLDQEPGPSQTPSILTSLGSTPIIEQSTKGGVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\hat{A}\mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.