

## 32-7630: Recombinant Mouse Stem Cell Factor/SCF/c-Kit Ligand

**Gene :** Kitlg  
**Gene ID :** 17311  
**Uniprot ID :** P20826

### Description

Source: E.coli.  
MW :18.4kD.

Recombinant Mouse Stem Cell Factor is produced by our E.coli expression system and the target gene encoding Lys26-Ala189 is expressed. Mouse stem cell factor (SCF), is the ligand for the receptor-type protein-tyrosine kinase KIT. It plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell development, migration and function, and in melanogenesis. KITLG/SCF binding can activate several signaling pathways. It also promotes phosphorylation of PIK3R1, which is the regulatory subunit of phosphatidylinositol 3-kinase, and subsequent activation of the kinase AKT1. KITLG/SCF and KIT also transmit signals via GRB2 and activation of RAS, RAF1 and the MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1. KITLG/SCF and KIT promote activation of STAT family members STAT1, STAT3 and STAT5.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** MKEICGNPVTDNVVDITKLVANLPNDYMITLNYVAGMDVLP SHCWLRDMVIQLSLSLTLLDKFSNIS  
EGLSNYSIIDKLGKIVDDL VLCMEENAPKNIKESPKRPETRSFTPEEFFSIFNRSIDAFKDFMVASDTS  
DCVLSSTLGPEKDSRVSVTKPFMLPPVA

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.