

32-7640: Recombinant Human CDK2AP2 (C-6His)

Gene : CDK2AP2

Gene ID : 10263

Uniprot ID : O75956

Description

Source: Human Cells.

MW :14.1kD.

Recombinant Human CDK2AP2 is produced by our Mammalian expression system and the target gene encoding Met1-Thr126 is expressed with a 6His tag at the C-terminus. CDK2AP2, also known as DOC1R, is short for cyclin-dependent kinase 2-associated protein 2. The gene CDK2AP2 encodes this protein that interacts with cyclin-dependent kinase 2 associated protein 1. Pseudogenes associated with this gene are located on chromosomes 7 and 9. Alternatively spliced transcript variants have been observed for this gene. It belongs to the CDK2AP family. CDK2AP1 (cyclin-dependent kinase 2-associated protein 1), corresponding to the gene doc-1 (deleted in oral cancer 1), is a tumor suppressor protein. The doc-1 gene is absent or down-regulated in hamster oral cancer cells and in many other cancer cell types. The ubiquitously expressed CDK2AP1 protein is the only known specific inhibitor of CDK2, making it an important component of cell cycle regulation during G(1)-to-S phase transition.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : MSYKPIAPAPSSTPGSSTPGPGTPVPTGSPSPSGSVPGAGAPFRPLFNDFGPPSMGYVQAMKP
PGAQGSQSTYTDLLSVIEEMGKEIRPTYAGSKSAMERLKRGIHARALVRECLAETERNARTVDHH
HHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.