

## 32-7655: Recombinant Human Serpin H1 (C-6His)

**Gene :** SERPINH1

**Gene ID :** 871

**Uniprot ID :** P50454

### Description

Source: Human Cells.

MW :45.5kD.

Recombinant Human Serpin H1 is produced by our Mammalian expression system and the target gene encoding Ala19-Leu418 is expressed with a 6His tag at the C-terminus. Serpin H1 is a serine proteinase inhibitors Which belongs to the serpin family. Serpin H1 is induced by heat shock. Serpin H1 localizes to the endoplasmic reticulum lumen and binds specifically to collagen. Thus it is thought to be a molecular chaperone involved in the maturation of collagen molecules. Autoantibodies to this protein have been found in patients with rheumatoid arthritis. Serpin H1 may be a marker for cancer and nucleotide polymorphisms in this gene may be associated with preterm birth caused by preterm premature rupture of membranes.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** AEVKKPAAAAAPGTAEKLSPKAATLAERSAGLAFSLYQAMAKDQAVENILVSPVWVASSLGLVSLG  
GKATTASQAKAVLSAEQLRDEEVHAGLGELLRSLSNSTARNVTWKLGSRLYGPSSVSFADDFVRS  
SKQHYNCEHSKINFRDKRSALQSINEWAAQTDDGKLPEVTKDVERTDGALLVNAMFFKPHWDEKF  
HHKMVDNRGFMVTRSYTVGVMMMHRTGLYNYDDKEKLQIVEMPLAHKLSSLILMPHHVEPLE  
RLEKLLTKEQLKIWMGKMQKKAIVSLPKGVVEVTHDLQKHLAAGLGLTEAIDKNKADLSRMSGKKD  
LYLASVFHATAFELDTDGNPFDQDIYGREELRSPKLFYADHPFIFLVRDTQSGSLLFIGRLVRPKGD  
KMRDELVDHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.