

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7656: Recombinant Human Myelin Protein P0/MPZ (C-6His)

**Gene :** MPZ **Gene ID :** 4359 **Uniprot ID :** P25189

## **Description**

Source: Human Cells.

MW:15.2kD.

Recombinant Human Myelin Protein P0 is produced by our Mammalian expression system and the target gene encoding Ile30-Arg153 is expressed with a 6His tag at the C-terminus. Myelin Protein P0 (MPZ) is a single-pass type I membrane glycoprotein which belongs to the myelin P0 protein family. MPZ contains one Ig-like V-type (immunoglobulin-like) domain, absent in the central nervous system. MPZ is a major component of the myelin sheath in peripheral nerves. It is postulated that MPZ is a structural element in the formation and stabilisation of peripheral nerve myelin, holding its characteristic coil structure together by the interaction of its positively-charged domain with acidic lipids in the cytoplasmic face of the opposed bilayer, and by interaction between hydrophobic globular of adjacent extracellular domains. Defects in MPZ associated with Charcot-Marie-Tooth disease and Dejerine-Sottas disease.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition:** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: IVVYTDREVHGAVGSRVTLHCSFWSSEWVSDDISFTWRYQPEGGRDAISIFHYAKGQPYIDEVGTF

KERIQWVGDPRWKDGSIVIHNLDYSDNGTFTCDVKNPPDIVGKTSQVTLYVFEKVPTRVDHHHHH

Η

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\hat{A}\mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.