

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7658: Recombinant Human Ribose-Phosphate Pyrophosphokinase 2/PRPS2 (C-6His)

Gene ID: 5634 **Uniprot ID:** P11908

Description

Source: Human Cells.

MW:35.8kD.

Recombinant Human PRPS2 is produced by our Mammalian expression system and the target gene encoding Pro2-Leu318 is expressed with a 6His tag at the C-terminus. Ribose-Phosphate Pyrophosphokinase 2 (PRPS2) is a phosphoribosyl pyrophosphate synthetase that belongs to the ribose-phosphate pyrophosphokinase family. PRPS2 is a homodimer. The active form is probably an hexamer composed of three homodimers. PRPS2 catalyzes the synthesis of phosphoribosylpyrophosphate (PRPP) that is essential for nucleotide synthesis. PRPS2 catalyzes the synthesis of 5-phosphoribosyl 1-pyrophosphate from ATP and D-ribose 5-phosphate. In addition, PRPS2 plays a central role in the synthesis of purines and pyrimidines.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: PNIVLFSGSSHQDLSQRVADRLGLELGKVVTKKFSNQETSVEIGESVRGEDVYIIQSGCGEINDNLMELLIMINAC

KIASSSRVTAVIPCFPYARQDKKDKSRAPISAKLVANMLSVAGADHIITMDLHASQIQGFFDIPVDNLYAEPAVLQ WIRENIAEWKNCIIVSPDAGGAKRVTSIADRLNVEFALIHKERKKANEVDRMVLVGDVKDRVAILVDDMADTCG TICHAADKLLSAGATKVYAILTHGIFSGPAISRINNAAFEAVVVTNTIPQEDKMKHCTKIQVIDISMILAEAIRRTHN

GESVSYLFSHVPLVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.