

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7661: Recombinant Human Galectin 9 (C-6His)(Discontinued)

Gene ID: LGALS9
Gene ID: 3965
Uniprot ID: Q3B8N1

## **Description**

Source: 293HEK cells.

MW:36.9kD.

Recombinant Human Galectin 9 is produced by our Mammalian expression system and the target gene encoding Met1-Thr323 is expressed with a 6His tag at the C-terminus. Galectin-9 is a cytoplasmic protein that contains two galectin domains. Galectin-9 is an S-type lectin that is over-expressed in Hodgkin's disease tissue. Galectin-9 binds galactosides and has high affinity for the Forssman pentasaccharide. Galectin-9 plays a role in thymocyte-epithelial interactions relevant to the biology of the thymus and Inhibits cell proliferation. Galectin-9 is a ligand for HAVCR2/TIM3 and induces T-helper type 1 lymphocyte (Th1) death. In addition, Galectin-9 suppresses tumor cell metastasis by interfering with the associations CD44, VCAM-1, Integrin a4 beta1.

## **Product Info**

Amount:  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of MOPs. 50mM NaCl, 1mM EDTA, 2mM DTT, pH 7.4

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

**Storage condition :** Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: MAFSGSQAPYLSPAVPFSGTIQGGLQDGLQITVNGTVLSSSGTRFAVNFQTGFSGNDIAFHFNPRF

EDGGYVVCNTRQNGSWGPEERKTHMPFQKGMPFDLCFLVQSSDFKVMVNGILFVQYFHRVPFHR VDTISVNGSVQLSYISFQPPGVWPANPAPITQTVIHTVQSAPGQMFSTPAIPPMMYPHPAYPMPFIT TILGGLYPSKSILLSGTVLPSAQRFHINLCSGNHIAFHLNPRFDENAVVRNTQIDNSWGSEERSLPRK

MPFVRGQSFSVWILCEAHCLKVAVDGQHLFEYYHRLRNLPTINRLEVGGDIQLTHVQT

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\mu$ g/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test.




Fig.1: Human Galectin-9, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.