

## 32-7662: Recombinant Human HVEM/TNFRSF14 (C-Fc)

**Gene :** TNFRSF14

**Gene ID :** 8764

**Uniprot ID :** Q92956

### Description

Source: Human Cells.

MW :44.5kD.

Recombinant Human Tumor Necrosis Factor Receptor Superfamily Member 14 is produced by our Mammalian expression system and the target gene encoding Leu39-Val202 is expressed with a Fc tag at the C-terminus. Herpesvirus entry mediator (HVEM) is a type I membrane protein in the TNF receptor superfamily, and it can both promote and inhibit T cell activity. HVEM is highly expressed on naïve CD4<sup>+</sup> T cells, CD8<sup>+</sup> T memory cells, regulatory T cells, dendritic cells, monocytes, and neutrophils. It functions as a receptor for BTLA, CD160, LIGHT/TNFSF14, and Lymphotoxin-α. Ligand of HVEM by LIGHT triggers T cell, monocyte, and neutrophil activation and contributes to Th1 inflammation and cardiac allograft rejection. In contrast, HVEM binding to CD160 or BTLA suppresses T cell and dendritic cell activation and dampens intestinal inflammation. HVEM enhances the development of CD8<sup>+</sup> T cell memory and Treg function. It is additionally expressed on intestinal epithelial cells, where its binding by intraepithelial lymphocyte (IEL) expressed CD160 promotes epithelial integrity and host defense. The herpesvirus envelope glycoprotein gD, which binds HVEM to initiate membrane fusion, can antagonize both BTLA and LIGHT binding.

### Product Info

**Amount :** 10 µg / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** LPSCKEDEYPVGSECCPKSPGYRVKEACGELTGTVCEPCPPGTYYIAHLNGLSKCLQCQMCDPAMGLRASRNC  
SRTENAVCGCSPGHFCIVQDGDHCAACRAYATSSPGQVRVQKGGTESQDTLCQNCPPGTFSFNGTLEECQHQT  
KCSWLVTGAGAGTSSSHWVDDIEGRMDPEKSCDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTC  
VVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIE  
KTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS  
KLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.