

32-7669: Recombinant Human Protamine-2/PRM2 (C-6His)

Gene : RRM2
Gene ID : 6241
Uniprot ID : P31350

Description

Source: Human Cells.
MW :45.9kD.

Recombinant Human Protamine-2/ is produced by our Mammalian expression system and the target gene encoding Met1-Phe389 is expressed with a 6His tag at the C-terminus. Ribonucleoside-Diphosphate Reductase Subunit M2 (RRM2) belongs to the ribonucleoside diphosphate reductase small chain family. The reductase of RRM2 catalyzes the formation of deoxyribonucleotides from ribonucleotides. Synthesis of the encoded protein (M2) is regulated in a cell-cycle dependent fashion. RRM2 supplies the precursors essential for DNA synthesis. RRM2 catalyzes the biosynthesis of deoxyribonucleotides from the corresponding ribonucleotides. Phosphorylation on Ser-20 relieves the inhibitory effect on Wnt signaling.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : MLSLRVPLAPITDPQQLQLSPLKGLSLVDKENTPPALSGTRVLASKTARRIFQEPTPEKTKAAAPGV
EDEPLLRENPRRFVIFPIEYHDIWQMYKKAEEASFWEVDLSKDIQHWESLKPEERYFISHVLAFF
AASDGIVNENLVERFSQEVQITEARCFYGFQIAMENIHSEMYSLIDTYIKDPKEREFLFNAIETMPCV
KKKADWALRWIGDKEATYGERVVFAAVEGIFFSGSFASIFWLKKRGLMPGLTFSNELISRDEGLH
CDFACLMTFKHLVHKPSEERVREIIINAVRIEQEFLTEALPVKLIGMNCTLMKQYIEFVADRLMLELGFS
KVFRVENPFDPMENISLEGKTNFFEKRVGEYQRMGMSSPTENSFTLDADFLEHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.