

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7675: Recombinant Human VIP36/LMAN2/GP36b (C-6His)

Gene ID: LMAN2
Gene ID: 10960
Uniprot ID: Q12907

Description

Source: Human Cells.

MW:32.7kD.

Recombinant Human Vesicular Integral-Membrane Protein VIP36 is produced by our Mammalian expression system and the target gene encoding Asp45-Arg322 is expressed with a 6His tag at the C-terminus. Vesicular integral-membrane protein VIP36 is also known as Glycoprotein GP36b, Lectin mannose-binding 2, Vesicular integral-membrane protein 36, LMAN2 and C5orf8. LMAN2 is widely expressed and contains one L-type lectin-like domain. LMAN2 binds high mannose type glycoproteins and may facilitate their sorting, trafficking and quality control. LMAN2 plays a role as an intracellular lectin in the early secretory pathway. LMAN2 interacts with N-acetyl-D-galactosamine and high-mannose type glycans and may also bind to O-linked glycans. LMAN2 is also involved in the transport and sorting of glycoproteins carrying high mannose-type glycans.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 50mM TrisHCl,10mM GSH,pH8.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: DITDGNSEHLKREHSLIKPYQGVGSSSMPLWDFQGSTMLTSQYVRLTPDERSKEGSIWNHQPCFL

KDWEMHVHFKVHGTGKKNLHGDGIALWYTRDRLVPGPVFGSKDNFHGLAIFLDTYPNDETTERVF PYISVMVNNGSLSYDHSKDGRWTELAGCTADFRNRDHDTFLAVRYSRGRLTVMTDLEDKNEWKN CIDITGVRLPTGYYFGASAGTGDLSDNHDIISMKLFQLMVEHTPDEESIDWTKIEPSVNFLKSPKDNV

DDPTGNFRSGPLTGWRVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\hat{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.