

32-7691: Recombinant Human Lactotransferrin/LTF (C-6His)

Gene : LTF
Gene ID : 4057
Uniprot ID : P02788

Description

Source: Human Cells.
MW :77.3kD.

Recombinant Human Lactotransferrin/LTF is produced by our Mammalian expression system and the target gene encoding Gly20-Lys710 is expressed with a 6His tag at the C-terminus. Lactotransferrin is a member of the transferrin family that transfer iron to the cells and control the level of free iron in the blood and external secretions. Lactotransferrin is a secreted protein and contains two transferrin-like domains. Lactotransferrin can be cleaved into the following four chains: Kaliocin-1, Lactoferroxin-A, Lactoferroxin-B, and Lactoferroxin-C. Lactoferroxin A, Lactoferroxin B, and Lactoferroxin C have opioid antagonist activity. Lactoferroxin A shows preference for mu-receptors, while Lactoferroxin B and Lactoferroxin C have somewhat higher degrees of preference for kappa-receptors than for mu-receptors. LTF has antimicrobial activity (bactericide, fungicide) and is part of the innate defense, mainly at mucoses.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : GSRRSVQWCAVSQPEATKCFQWQRNMRKVRGPPVSCIKRDSPIQCIQAIENRADAVTLDGGFIYEAGLAPY KLRPVAAEVYGTQRPRTHYYAVAVVKKGGSFQLNELQGLKSCHTGLRRTAGWNVPIGTLRPFLNWTGPPEPIE AAVARFFSASCVPADKGQFPNLCRLCAGTGGENKCAFSSQEPYFSYSGAFKCLRDGAGDVAFIRESTVFEDLS DEARDEYELLCPDNTRKPVDFKDKCHLARVPASHVARSVNGKEDAIWNLLRQAQEKFGKDKSPKFQFLGSP SGQKDLLFKDSAIGFSRVPPRIDSGLYLGGYFTAIQNLKSEEEVAARRARVVWCAVGEQELRKCQNQWSGLS EGSVTCSSASTTEDICIALVLKGEADAMSLDGGYVYTAGKCGLPVLAENYKSQQSSDPDPNCVDRPVEGYLTV AVVRRSDTSLTWNSVKGKKSCHTAVDRTAGWNIPMGLLFNQTGSKCFDEYFSQSCAPGSDPRSNLCALCIGD EQGENKCEPNSNERYGYTGAFRCLAENAGDVAFVKDVTVLQNTDGNNEAWAKDLKLADFALLCLDGKRPK VTEARSCHLAMAPNHAVVSMDKVERLKQVLLHQAKFGRNGSDCPDKFCLFQSETKNLLFNDNTECLARLH GKTTYEKYLGPPYVAGITNLKCKSTSPLEACEFLRKVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.