9853 Pacific Heights Blvd.
Suite D. San Diego,
CA 92121, USA
Tel: 858-263-4982
Email: info@abeomics.com

32-7702: Recombinant Human CEACAM7/CGM2 (C-6His)

Gene : CEACAM7
Gene ID : 1087
Uniprot ID : Q14002

Description

Source: Human Cells.
MW :13.1kD.
Recombinant Human CEACAM7 is produced by our Mammalian expression system and the target gene encoding Thr36-Phe142 is expressed with a 6His tag at the C-terminus. Carcinoembryonic Antigen-Related Cell Adhesion Molecule 7 (CEACAM7) is a member of the immunoglobulin superfamily A and CEA family. CEACAM7 localizes to the cell membrane and contains one Ig-like C2-type domain and one Ig-like V-type domain. The expression of CEACAM7 is significantly decreased in rectal cancer. Differences in CEACAM7 expression levels between long-term survivors and those with recurrent disease introduce a potential tumor marker to define a subset of patients who benefit most from adjuvant therapy.

Product Info

Amount :	10 µg / 50 µg
Content :	Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition :	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid :	TNIDVVPFNVAEGKEVLLVVHNESQNLYGYNWYKGERVHANYRIIGYVKNISQENAPGPAHNGRETIYPNGTLLI QNVTHNDAGIYTLHVIKENLVNEEVTRQFYVFVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Ã‚Âµg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Endotoxin : Less than 0.1 ng/Ã‚Âµg (1 IEU/Ã‚Âµg) as determined by LAL test.

For Research Use Only. Not for use in diagnostic/therapeutics procedures.