

32-7705: Recombinant Human PACSIN2/Syndapin-2 (C-6His)(Discontinued)

Gene : PACSIN2

Gene ID : 11252

Uniprot ID : Q9UNF0

Description

Source: Human Cells.

MW :56.7kD.

Recombinant Human Syndapin-2 is produced by our Mammalian expression system and the target gene encoding Met1-Gln486 is expressed with a 6His tag at the C-terminus. Protein Kinase C and Casein Kinase Substrate in Neurons Protein 2 (PACSIN2) is a member of the PACSIN family. PACSIN2 is localized to the plasma membrane via its coiled-coil domain. PACSIN2 is widely expressed and contains one FCH domain and one SH3 domain. PACSIN2 forms homo- and hetero-aggregates with other PACSINs. PACSIN2 may play a role in vesicle formation and transport. In addition, PACSIN2 is involved in linking the actin cytoskeleton with vesicle formation by regulating tubulin polymerization.

Product Info

Amount : 10 µg / 50 µg

Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

Amino Acid : MSVTYDDSVGVEVSSDSFWEVGNKYKRTVKRIDDGHRCLSDLMNCLHERARIEKAYAQQLTEWARRWRQLVE KGPQYGTVEKAWMAFMSEAERVSELHLEVKASLMNDDFEKIKNWQKEAFHKQMMGGFKETKEAEDGFRKAQ KPWAKKLKEVEAAKKAHHAACKEEKLAISREANSKADPSLNPEQLKKLQDKIECKQDVLKTKKEYEKSLELD QGTPQYMNMEQVFEQCQQFEEKRLRFFREVLLVQKHLDSLNVAGYKAIYHDLEQSIRAADAVEDLRWFRAN HGPGMAMNWPQFEEWSADLNRTLRSRREKKKATDGVTLTGINQTDGQSLPSKPSSTLNVPSNPAQSAQSQSSY NPFEDDDTGSTVSEKDDTKAKNVSSYEKTQSYPTDWSDDSNPFSSTDANGDSNPFDDDATSGTEVRVRA LYDYEGQEHDELSFKAGDELTKMEDEDEQGWCKGRDLNGQVGLYPANYVEAIQVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.