

## 32-7716: Recombinant Human Porphobilinogen Deaminase/HMBS/PBGD (C-6His)

**Gene :** HMBS  
**Gene ID :** 3145  
**Uniprot ID :** P08397

### Description

Source: Human Cells.

MW :40.5kD.

Recombinant Human Porphobilinogen Deaminase is produced by our Mammalian expression system and the target gene encoding Ser2-His361 is expressed with a 6His tag at the C-terminus. Porphobilinogen Deaminase (HMBS) is a member of the HMBS family. PBGD is the third enzyme of the heme biosynthetic pathway and catalyzes the head to tail condensation of four porphobilinogen molecules into the linear hydroxymethylbilane. HMBS is involved in the production of heme, which is important for all of the body's organs, although it is most abundant in the blood, bone marrow, and liver. In addition, Heme is an essential component of iron-containing proteins called hemoproteins, including hemoglobin. Defects in PBGD are the cause of acute intermittent porphyria.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** SGNGNAAATAEENSPKMRVIRVGTRKSQLARIQTDSVVATLKASYPGLQFEIIMSTTGDKILDALTALSKIGEKSLF  
TKELEHALEKNEVDLVVHSLKDLPTVLPFGFTIGAICKRENPHDAVVFHPKFVGKTLETLPKSVVGTSSLRRAA  
QLQRKFPHEFRSIRGNLNTLRKLDEQQEFSAILATAGLQRMGWHNRVGQILHPEECMYAVGQGALGVEVR  
AKDQDILDVLGVLDHPETLLRCIAERAFRLRHLEGGCSVPVAVHTAMKDGQLYLTGGVWSLDGSDSIQETMQAT  
IHVPAQHEDGPEDDPQLVGITARNIPRGPQLAAQNLGISLANLLLSKGAKNILDVARQLNDAHVDHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.