

32-7720: Recombinant Human Vitamin D-Binding Protein/VDB/Gc-globulin (C-6His)

Gene : GC
Gene ID : 2638
Uniprot ID : P02774

Description

Source: Human Cells.
MW :52.3kD.

Recombinant Human Vitamin D-Binding Protein is produced by our Mammalian expression system and the target gene encoding Leu17-Leu474 is expressed with a 6His tag at the C-terminus. Vitamin D-Binding Protein (DBP) is a member of the ALB/AFP/VDB family. DBP is a secreted protein and contains three albumin domains. The primary structure contains 28 cysteine residues forming multiple disulfide bonds. DBP acts as a multifunctional protein found in plasma, ascitic fluid, cerebrospinal fluid, and urine and on the surface of many cell types. DBP binds to vitamin D and its plasma metabolites and transports them to target tissues. DBP associates with membrane-bound immunoglobulin on the surface of B-lymphocytes and with IgG Fc receptor on the membranes of T-lymphocytes.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.2.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : LERGRDYEKNKVCKEFSLGKEDFTSLSLVLYSRKFPSGTFEQVSQLVKEVVSLTEACCAEGADPDCYDTRTSA
LSAKSCESNSPPFVHPGTAECCTKEGLERKLCMAALKHQPEFPTYVEPTNDEICEAFRKDPKEYANQFMWEYS
TNYGQAPLSLLVSYTKSYLSMVGSCCTSASPTVCFLKERLQLKHLSTLTLNVRVCSQYAAAYGEKKSRLSNLIKLA
QKVPTADLEDVLPLAEDITNILSKCCESASEDCMAKELPEHTVKLCDNLSTKNSKFEDCCQEKAMDFVCTYF
MPAAQLPELPDVELPTNKDVCDPGNTKVMKYTFELSRRLHLPEVFLSKVLEPTLKSLECCDVEDSTTCFNAK
GPLLKKELSSFIDKGQELCADYSENTFTYKKLAERLAKLPDATPTLAKLVNKRSDFASNCCSINSPLYCDS
EIDAELKNILVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.