

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7725: Recombinant Human B3GNT1 (C-6His)(Discontinued)

Gene ID: 84GAT1 **Gene ID:** 11041 **Uniprot ID:** 043505

Description

Source: Human Cells.

MW:43.4kD.

Recombinant Human B3GNT1 is produced by our Mammalian expression system and the target gene encoding Asp43-Cys415 is expressed with a 6His tag at the C-terminus. N-Acetyllactosaminide beta-1,3-N-Acetylglucosaminyltransferase (B3GNT1) is a member of the beta-1,3-N-Acetylglucosaminyltransferase family. B3GNT1 is a single-pass type II membrane protein and widely expressed in many tissues. B3GNT1 can initiate the synthesis or the elongation of the linear poly-N-acetyllactosaminoglycans. B3GNT1 is essential for the synthesis of poly-N-acetyllactosamine, a determinant for the blood group i antigen. It can initiate the synthesis or the elongation of the linear poly-N-acetyllactosaminoglycans.

Product Info

Amount: 10 μg / 50 μg

Content : Lyophilized from a 0.2 µm filtered solution of PBS,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: DQYFEFFPPSPRSVDQVKAQLRTALASGGVLDASGDYRVYRGLLKTTMDPNDVILATHASVDNLLHLSGLLER

WEGPLSVSVFAATKEEAQLATVLAYALSSHCPDMRARVAMHLVCPSRYEAAVPDPREPGEFALLRSCQEVFDK LARVAQPGINYALGTNVSYPNNLLRNLAREGANYALVIDVDMVPSEGLWRGLREMLDQSNQWGGTALVVPAF EIRRARRMPMNKNELVQLYQVGEVRPFYYGLCTPCQAPTNYSRWVNLPEESLLRPAYVVPWQDPWEPFYVAG GKVPTFDERFRQYGFNRISQACELHVAGFDFEVLNEGFLVHKGFKEALKFHPQKEAENQHNKILYRQFKQELKA

KYPNSPRRCVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \Box \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.