

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7727: Recombinant Human Cysteine-Rich with EGF-Like Domain Protein 2/CRELD2 (C-6His)(Discontinued)

CRELD2 Gene: Gene ID: 79174 Uniprot ID: Q6UXH1

## **Description**

Source: Human Cells.

MW:33.4kD.

Recombinant Human CRELD2 is produced by our Mammalian expression system and the target gene encoding Ala25-Leu321 is expressed with a 6His tag at the C-terminus. Cysteine-Rich with EGF-Like Domain Protein 2 (CRELD2) is a secreted protein that is a member of the CRELD family. Human CRELD2 is synthesized as a 353 amino acid precursor protein with a signal peptide, a highly conserved domain rich in glutamic acid and tryptophan (WE) and EGF-like repeats. CRELD2 is ubiquitously expressed in many tissues. CRELD2 may interact with CHRNA4 and regulate transport of a4- beta2 neuronal acetylcholine receptor. In addition, CRELD2 could be a novel mediator in regulating the onset and progression of various ER stressassociated diseases.

## **Product Info**

Amount:  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of PBS,5%Trehalose, pH 7.4.

> Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

AKKPTPCHRCRGLVDKFNQGMVDTAKKNFGGGNTAWEEKTLSKYESSEIRLLEILEGLCESSDFEC **Amino Acid:** 

NQMLEAQEEHLEAWWLQLKSEYPDLFEWFCVKTLKVCCSPGTYGPDCLACQGGSQRPCSGNGH CSGDGSRQGDGSCRCHMGYQGPLCTDCMDGYFSSLRNETHSICTACDESCKTCSGLTNRDCGE CEVGWVLDEGACVDVDECAAEPPPCSAAQFCKNANGSYTCEDVDECSLAEKTCVRKNENCYNTP

GSYVCVCPDGFEETEDACVPPAEAEATEGESPTQLPSREDLVDHHHHHH

## **Application Note**

Storage condition:

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.