

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7741: Recombinant Human Apolipoprotein A-IV/ApoA4 (C-6His)

Gene: APOA4 Uniprot ID: P06727

## **Description**

Source: Human Cells. MW:44.44kD.

Recombinant Human Apolipoprotein A-IV is produced by our Mammalian expression system and the target gene encoding Glu21-Ser396 is expressed with a 6His tag at the C-terminus. Apolipoprotein A4 (APOA4) is a secreted protein that belongs to the apolipoprotein A1/A4/E family. Apoa-IV is a major component of HDL and chylomicrons. APOA4 is secreted into circulation on the surface of newly synthesized chylomicron particles. APOA4 play a role in the regulation of appetite and satiety in rodent models. APOA4 involved in chylomicrons and VLDL secretion and catabolism and required for efficient activation of lipoprotein lipase by ApoC-II. In addition, APOA4 is a potent activator of lecithin-cholesterol acyltransferase in vitro.

## **Product Info**

Amount:  $10 \mu g / 50 \mu g$ 

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition:

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: EVSADQVATVMWDYFSQLSNNAKEAVEHLQKSELTQQLNALFQDKLGEVNTYAGDLQKKLVPFAT

ELHERLAKDSEKLKEEIGKELEELRARLLPHANEVSQKIGDNLRELQQRLEPYADQLRTQVNTQAE QLRRQLTPYAQRMERVLRENADSLQASLRPHADELKAKIDQNVEELKGRLTPYADEFKVKIDQTVE ELRRSLAPYAQDTQEKLNHQLEGLTFQMKKNAEELKARISASAEELRQRLAPLAEDVRGNLRGNTE GLQKSLAELGGHLDQQVEEFRRRVEPYGENFNKALVQQMEQLRQKLGPHAGDVEGHLSFLEKDL

RDKVNSFFSTFKEKESQDKTLSLPELEQQQEQQQEQQQEQVQMLAPLESVDHHHHHH

## **Application Note**

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\hat{A}\mu g/ml$ . Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin**: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.