

## 32-7751: Recombinant Human Fibroblast Growth Factor 23/FGF-23 (C-6His)(Discontinued)

**Gene :** FGF23  
**Gene ID :** 8074  
**Uniprot ID :** Q9GZV9

### Description

Source: Human Cells.  
MW :26.4kD.

Recombinant Human Fibroblast Growth Factor 23 is produced by our Mammalian expression system and the target gene encoding Tyr25-Ile251 is expressed with a 6His tag at the C-terminus. Fibroblast Growth Factor 23 (FGF-23) is a secreted protein that belongs to the heparin-binding growth factors family. FGF-23 is expressed in osteogenic cells, particularly during phases of active bone remodeling. FGF family members possess broad mitogenic and cell survival activities, involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth, and invasion. FGF-23 regulates homeostasis of phosphate and vitamin-D metabolism. FGF-23 inhibits renal tubular phosphate transport by reducing SLC34A1 levels, and negatively regulates osteoblast differentiation and matrix mineralization. FGF-23 also upregulates EGR1 expression in the presence of KL, acts directly on the parathyroid to decrease PTH secretion. Defects in FGF-23 are the cause of autosomal dominant hypophosphataemic rickets (ADHR).

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,1mM EDTA,2mMDTT,pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** YPNASPLLGSWGGLIHLYTATARN SYHLQIHKN GHVDGAPHQTIYSALMIRSEDAGFVVITGVMSRRYLCMDF  
RGNIFGSHYFDPENCRFQHQTLENGYDVYHSPQYHFLVSLGRAKRAFLPGMNPPYPYSQFLSRRNEIPLIHFNTP  
IPRRHTRSAEDDSERDPLNVLKPRARMT PAPASCSQELPSAEDNSPMASDPLGVVRGGRVNT HAGGTGPEGCR  
PFAKFIVDHHHHHHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.