

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7754: Recombinant Human GM-CSF R a/CSF2RA/CD116 (C-6His)

Gene ID: CSF2RA
Gene ID: 1438
Uniprot ID: P15509

Description

Source: Human Cells.

MW:35.5kD.

Recombinant Human GM-CSF Receptor alpha is produced by our Mammalian expression system and the target gene encoding Glu23-Gly320 is expressed with a 6His tag at the C-terminus. Granulocyte-Macrophage Colony-Stimulating Factor Receptor Subunit a (CSF2RA) is a single-pass type I membrane protein which belongs to the type I cytokine receptor family of Type 5 subfamily. The CSF2RA gene is found in the pseudoautosomal region (PAR) of the X and Y chromosomes with some of the isoforms being membrane-bound and others being soluble. CSF2RA is a low affinity receptor for granulocyte-macrophage colony-stimulating factor. CSF2RA transduces a signal that results in the proliferation, differentiation, and functional activation of hematopoietic cells. Defects in CSF2RA are the cause of pulmonary surfactant metabolism dysfunction type 4 (SMDP4).

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: EKSDLRTVAPASSLNVRFDSRTMNLSWDCQENTTFSKCFLTDKKNRVVEPRLSNNECSCTFREICLHEGVTFEV

HVNTSQRGFQQKLLYPNSGREGTAAQNFSCFIYNADLMNCTWARGPTAPRDVQYFLYIRNSKRRREIRCPYYIQ DSGTHVGCHLDNLSGLTSRNYFLVNGTSREIGIQFFDSLLDTKKIERFNPPSNVTVRCNTTHCLVRWKQPRTYQ KLSYLDFQYQLDVHRKNTQPGTENLLINVSGDLENRYNFPSSEPRAKHSVKIRAADVRILNWSSWSEAIEFGSD

DGVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 $ng/\mu g$ (1 $IEU/\mu g$) as determined by LAL test.