

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7770: Recombinant Human Insulin-Like Growth Factor-Binding Protein 5/IGFBP-5 (C-6His)

Gene ID: 3488
Uniprot ID: P24593

Description

Source: Human Cells. MW:29.61kD.

Recombinant Human Insulin-Like Growth Factor-Binding Protein 5 is produced by our Mammalian expression system and the target gene encoding Leu21-Glu272 is expressed with a 6His tag at the C-terminus. Insulin-Like Growth Factor-Binding Protein 5 (IGFBP-5) is a secreted protein that belongs to the insulin-like growth factor (IGF) binding proteins superfamily. Members of this family prolong the half-life of the IGFs. They have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors. IGFBP-5 contains one IGFBP N-terminal domain and one thyroglobulin type-1 domain. IGFBP-5 is expressed by fibroblasts, myoblasts and Osteosarcoma. It is also present at lower levels in liver, kidney, and brain.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: LGSFVHCEPCDEKALSMCPPSPLGCELVKEPGCGCCMTCALAEGQSCGVYTERCAQGLRCLPRQDEEKPLHAL

LHGRGVCLNEKSYREQVKIERDSREHEEPTTSEMAEETYSPKIFRPKHTRISELKAEAVKKDRRKKLTQSKFVGG AENTAHPRIISAPEMRQESEQGPCRRHMEASLQELKASPRMVPRAVYLPNCDRKGFYKRKQCKPSRGRKRGIC

WCVDKYGMKLPGMEYVDGDFQCHTFDSSNVEVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than $0.1 \text{ ng}/\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.