

32-7772: Recombinant Human Calumenin/CALU (C-6His)

 Gene :
 CALU

 Gene ID :
 813

 Uniprot ID :
 043852

Description

Source: Human Cells. MW :36kD.

Recombinant Human Calumenin is produced by our Mammalian expression system and the target gene encoding Lys20-Phe315 is expressed with a 6His tag at the C-terminus. Calumenin is a secreted calcium-binding protein that belongs to the CREC family. Calumenin contains six EF-hand domains and is expressed at high levels in the heart, placenta and skeletal muscle. Human Calumenin is synthesized as a 315 amino acid precursor that contains a 19 amino acid signal sequence, and a 296 amino acid mature chain. Calumenin localizes to the endoplasmic reticulum (ER) and sarcoplasmic reticulum (SR) of mammalian tissues which plays a role in ER functions as protein folding and sorting. Calumenin is involved in the regulation of vitamin K-dependent carboxylation of multiple N-terminal glutamate residues. It seems to inhibit gamma -carboxylase GGCX.

Product Info

Amount : Content :	10 μg / 50 μg Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition :	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid :	KPTEKKDRVHHEPQLSDKVHNDAQSFDYDHDAFLGAEEAKTFDQLTPEESKERLGKIVSKIDGDKDGFVTVDE LKDWIKFAQKRWIYEDVERQWKGHDLNEDGLVSWEEYKNATYGYVLDDPDPDDGFNYKQMMVRDERRFKM ADKDGDLIATKEEFTAFLHPEEYDYMKDIVVQETMEDIDKNADGFIDLEEYIGDMYSHDGNTDEPEWVKTEREQ FVEFRDKNRDGKMDKEETKDWILPSDYDHAEAEARHLVYESDQNKDGKLTKEEIVDKYDLFVGSQATDFGEAL VRHDEFVDHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 \tilde{A} $\tilde{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ \tilde{A}] $\hat{A}\mu g$ (1 IEU/ \tilde{A}] $\hat{A}\mu g$) as determined by LAL test.