

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7789: Recombinant Human Endoplasmic Reticulum Resident Protein 27/ERP27 (C-6His)

Gene : ERP27 Gene ID : 121506 Uniprot ID : Q96DN0

Description

Source: Human Cells. MW:28.79kD.

Recombinant Human ERP27 is produced by our Mammalian expression system and the target gene encoding Glu26-Leu273 is expressed with a 6His tag at the C-terminus. Endoplasmic reticulum resident protein 27, also known as ER protein 27, C12orf46 and ERP27, is an endoplasmic reticulum luminal protein which is a member of the protein disulfide isomerase family. ERP27 contains one thioredoxin domain and does not contain a CXXC active site motif. ERP27 is widely expressed in many tissues; it has highest expression in pancreas, with lower levels in spleen, lung, kidney, thymus, and bone marrow. ERP27 interacts with PDIA3 and binds somatostatin-14 via hydrophobic interactions. ERP27 may act as a protease, protein disulfide isomerase, thiol-disulfide oxidase or phospholipase.

Product Info

Amount: $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: EVEKSSDGPGAAQEPTWLTDVPAAMEFIAATEVAVIGFFQDLEIPAVPILHSMVQKFPGVSFGISTD

SEVLTHYNITGNTICLFRLVDNEQLNLEDEDIESIDATKLSRFIEINSLHMVTEYNPVTVIGLFNSVIQIH LLLIMNKASPEYEENMHRYQKAAKLFQGKILFILVDSGMKENGKVISFFKLKESQLPALAIYQTLDDE

WDTLPTAEVSVEHVQNFCDGFLSGKLLKENRESEGKTPKVELVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\hat{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.