

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7810: Recombinant Human Lecithin-cholesterol acyltransferase/LCAT (C-6His)

Gene ID: 3931
Uniprot ID: P04180

Description

Source: Human Cells.

MW:48.1kD.

Recombinant Human Lecithin-cholesterol acyltransferase is produced by our Mammalian expression system and the target gene encoding Phe25-Glu440 is expressed with a 6His tag at the C-terminus. Lipase family. The gene encoding this protein is expressed mainly in brain, liver and testes, followed by secreting into plasma and cerebral spinal fluid. The esterification of cholesterol is required for cholesterol transport. LCAT is a central enzyme in the extracellular metabolism of plasma lipoproteins. Defects in LCAT are the cause of lecithin-cholesterol acyltransferase deficiency (LCATD) and a cause of fish-eye disease (FED).

Product Info

Amount: 10 μg / 50 μg

Content : Lyophilized from a 0.2 µm filtered solution of 50mM Acetate Buffer pH-4.0.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Storage condition : Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: FWLLNVLFPPHTTPKAELSNHTRPVILVPGCLGNQLEAKLDKPDVVNWMCYRKTEDFFTIWLDLNMFLPLGVDC

WIDNTRVVYNRSSGLVSNAPGVQIRVPGFGKTYSVEYLDSSKLAGYLHTLVQNLVNNGYVRDETVRAAPYDWR LEPGQQEEYYRKLAGLVEEMHAAYGKPVFLIGHSLGCLHLLYFLLRQPQAWKDRFIDGFISLGAPWGGSIKPML VLASGDNQGIPIMSSIKLKEEQRITTTSPWMFPSRMAWPEDHVFISTPSFNYTGRDFQRFFADLHFEEGWYMWL OSRDLLAGLPAPGVEVYCLYGVGLPTPRTYIYDHGFPYTDPVGVLYEDGDDTVATRSTELCGLWQGRQPQPVH

LLPLHGIQHLNMVFSNLTLEHINAILLGAYRQGPPASPTASPEPPPPEVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\tilde{A} \square \hat{A} \mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/ $\tilde{A} \square \hat{A} \mu g$ (1 IEU/ $\tilde{A} \square \hat{A} \mu g$) as determined by LAL test.