

## 32-7830: Recombinant Mouse Folate Receptor Alpha/FOLR1 (C-6His)

**Gene :** Folr1  
**Gene ID :** 14275  
**Uniprot ID :** P35846

### Description

Source: Human Cells.  
MW :25.3kD.

Recombinant Mouse Folate Receptor Alpha is produced by our Mammalian expression system and the target gene encoding Thr25-Ser232 is expressed with a 6His tag at the C-terminus. Folate Receptor alpha belongs to the folate receptor family and it is a 37 - 42 kDa protein that mediates the cellular uptake of folic acid and reduced folates.. Mature FOLR1 is an N-glycosylated protein that is anchored to the cell surface by a GPI linkage. FOLR1 can be detected in kidney proximal tubules. It is critically required during early embryogenesis as shown in knockout mice which die in utero with gross morphological defects. FOLR1 binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells. It Has high affinity for folate and folic acid analogs at neutral pH. Exposure to slightly acidic pH after receptor endocytosis triggers a conformation change that strongly reduces its affinity for folates and mediates their release. Required for normal embryonic development and normal cell proliferation. Required for renal folate reabsorption.

### Product Info

**Amount :** 10 µg / 50 µg  
**Content :** Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.  
**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.  
**Amino Acid :** TRARTELLNVCMMDAKHHKEKPGPEDNLHDQCSPWKTNCCSTNTSQAHKDISYLYRFNWNHCGTMTSECK  
RHFIQDTCLEYECSPNLGPWIQQVDQSWRKERILDVPLCKEDCQQWVEDCQSSFTCKSNWHKGNWSSGHN  
ECPVGASCHPFTFYFPTSAALCEEIWSHSYKLSNYSRGSGRCIQMWFDPAGNPNEEVARFYAEAMSVDHHH  
HHH

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.