

32-7908: Recombinant Human Butyrophilin 3A3/BTN3A3 (C-6His)

 Gene :
 BTN3A3

 Gene ID :
 10384

 Uniprot ID :
 000478

Description

Source: Human Cells.

MW :24.6kD.

Recombinant Human Butyrophilin 3A3 is produced by our Mammalian expression system and the target gene encoding Gln30-Trp248 is expressed with a 6His tag at the C-terminus. Human BTN3A3, also known as butyrophilin subfamily 3 member A3 and BTF3, is a Single-pass type I membrane protein which belongs to the immunoglobulin superfamily and BTN/MOG family. The butyrophilin (BTN) genes are a group of major histocompatibility complex (MHC)-associated genes that encode type I membrane proteins with 2 extracellular immunoglobulin domains and an intracellular B30.2 (PRYSPRY) domain. It can be detected in peripheral blood mononuclear cells, T-cells I, spleen and lymphocytes. BTN3A3 plays a role in T-cell responses in the adaptive immune response.

Product Info

Amount :	10 µg / 50 µg
Content :	Lyophilized from a 0.2 μ m filtered solution of 20mM PB,150mM NaCl,pH7.4.
Storage condition :	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid :	QFSVLGPSGPILAMVGEDADLPCHLFPTMSAETMELRWVSSSLRQVVNVYADGKEVEDRQSAPYRGRTSILRD GITAGKAALRIHNVTASDSGKYLCYFQDGDFYEKALVELKVAALGSDLHIEVKGYEDGGIHLECRSTGWYPQPQI KWSDTKGENIPAVEAPVVADGVGLYAVAASVIMRGSSGGGVSCIIRNSLLGLEKTASISIADPFFRSAQPWVDH HHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 \tilde{A} $\hat{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/ \tilde{A} \square $\hat{A}\mu$ g (1 IEU/ \tilde{A} \square $\hat{A}\mu$ g) as determined by LAL test.