

32-7917: Recombinant Human IL-21 Receptor/IL-21R (C-6His)(Discontinued)

Gene : IL21R
Gene ID : 50615
Uniprot ID : Q9HBE5

Description

Source: Human Cells.
MW :26kD.

Recombinant Human Interleukin-21 Receptor is produced by our Mammalian expression system and the target gene encoding Cys20-Pro236 is expressed with a 6His tag at the C-terminus. Interleukin-21 receptor is also known as IL-21 receptor, IL-21R, CD360. In humans, it is encoded by the IL21R gene. It belongs to the type I cytokine receptor family. Type 4 subfamily contains 2 fibronectin type-III domains. Interleukin-21 receptor is selectively expressed in lymphoid tissues and highly expressed in thymus and spleen. IL-21 is produced by CD4+ T cells in response to antigenic stimulation. Its action enhances antigen-specific responses of immune cells. The biological effects of IL-21 include induction of differentiation of T-cells-stimulated B-cells into plasma cells and memory B-cells. It also promotes the anti-tumor activity of CD8+ T-cells and NK cells.

Product Info

Amount : 10 µg / 50 µg
Content : Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl, pH 7.4.
Storage condition : Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.
Amino Acid : CPDLVCYTDYLTQVICILEMWNLHPSTLTLTWQDQYEELKDEATSCSLHRSAHNATHATYTTCHMDV
FHFMAADDIFSVNITDQSGNYSQECGSFLAESIKPAPPFNVTVTFSGQYNISWRSDYEDPAFYMLK
GKLQYELQYRNRGDPWAVSPRRKLISVDSRSVLLPLEFRKDSSYELQVRAGPMPGSSYQGTWS
EWSDPVIFQTQSEELKEGWNPVDHHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin : Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.