

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

32-7923: Recombinant Human CEACAM21 (C-6His)

Gene : CEACAM21 **Gene ID :** 90273 **Uniprot ID :** Q3KPI0

Description

Source: Human Cells.

MW:24.2kD.

Recombinant Human CEACAM21 is produced by our Mammalian expression system and the target gene encoding Trp35-Gly240 is expressed with a 6His tag at the C-terminus. Carcinoembryonic antigen-related cell adhesion molecule 21 is a protein that in humans is encoded by the CEACAM21 gene. It belongs to the immunoglobulin superfamily. CEA family.containing 1 Iglike C2-type (immunoglobulin-like) domain. It was found to be a cell-cell adhesion molecule detected on leukocytes, epithelia, and endothelia. The encoded protein mediates cell adhesion via homophilic as well as heterophilic binding to other proteins of the subgroup. Multiple cellular activities have been attributed to the encoded protein, including roles in the differentiation and arrangement of tissue three-dimensional structure, angiogenesis, apoptosis, tumor suppression, metastasis, and the modulation of innate and adaptive immune responses.

Product Info

Amount : $10 \mu g / 50 \mu g$

Content: Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl, pH 7.4.

Lyophilized protein should be stored at -20 $^{\circ}\text{C},$ though stable at room temperature for 3 weeks.

Storage condition: Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: WLFIASAPFEVAEGENVHLSVVYLPENLYSYGWYKGKTVEPNQLIAAYVIDTHVRTPGPAYSGRETI

SPSGDLHFQNVTLEDTGYYNLQVTYRNSQIEQASHHLRVYESVAQPSIQASSTTVTEKGSVVLTCH TNNTGTSFQWIFNNQRLQVTKRMKLSWFNHVLTIDPIRQEDAGEYQCEVSNPVSSNRSDPLKLTVK

SDDNTLGVDHHHHHH

Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\hat{A}\mu g/ml$. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Endotoxin: Less than 0.1 ng/Âμg (1 IEU/Âμg) as determined by LAL test.