

9853 Pacific Heights Blvd. Suite D. San Diego, CA 92121, USA Tel: 858-263-4982

Email: info@abeomics.com

## 32-7933: Recombinant Human CTHRC1/NMTC1 (C-6His) in 293HEK cells

Gene ID: CTHRC1
Gene ID: 115908
Uniprot ID: Q96CG8

## **Description**

Source: Human Cells.

MW:24.1kD.

Recombinant Human CTHRC1 in 293 HEK mammalian expression system and the target gene encoding Ser31-Lys243 is expressed with a 6His tag at the C-terminus. Collagen triple helix repeat-containing protein 1 is a protein that in humans is encoded by the CTHRC1 gene. It acts as a negative regulator of collagen matrix deposition. It may cause the disease of Barrett esophagus . Patients with Barrett esophagus have an increased risk of esophageal adenocarcinoma. The main cause of Barrett esophagus is gastroesophageal reflux. The retrograde movement of acid and bile salts from the stomach into the esophagus causes prolonged injury to the esophageal epithelium and induces chronic esophagitis, which in turn is believed to trigger the pathologic changes.

## **Product Info**

**Amount :**  $10 \mu g / 50 \mu g$ 

**Content:** Lyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl,pH7.4.

Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted

samples are stable at -20°C for 3 months.

Amino Acid: SEIPKGKQKAQLRQREVVDLYNGMCLQGPAGVPGRDGSPGANGIPGTPGIPGRDGFKGEKGECLRESFEESW

TPNYKQCSWSSLNYGIDLGKIAECTFTKMRSNSALRVLFSGSLRLKCRNACCQRWYFTFNGAECSGPLPIEAIIYL DQGSPEMNSTINIHRTSSVEGLCEGIGAGLVDVAIWVGTCSDYPKGDASTGWNSVSRIIIEELPKVDHHHHHH

## **Application Note**

Storage condition:

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100  $\mu$ g/ml. Dissolve the lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than  $0.1 \text{ ng/}\mu\text{g}$  (1 IEU/ $\mu\text{g}$ ) as determined by LAL test.




Fig 1: SDS-PAGE analysis of Recombinant Human CTHRC1/NMTC1 (C-6His) 293 HEK cells. R lane is in reducing conditions, NR for non-reducing.