

## 32-7944: Recombinant Human Ephrin B Receptor 1/EphB1 (C-Fc)(Discontinued)

**Gene :** EPHB1

**Gene ID :** 2047

**Uniprot ID :** P54762

### Description

Source: Human Cells.

MW :85.5kD.

Recombinant Human Ephrin B Receptor 1 is produced by our Mammalian expression system and the target gene encoding Met18-Pro540 is expressed with a Fc tag at the C-terminus. Ephrin Type-B Receptor 1 (EPHB1) is a single-pass type I membrane protein that belongs to the Ephrin-B family of receptor tyrosine kinases involved in the development of embryonic nervous and vascular systems. EPHB1 contains two fibronectin type-III domains, one protein kinase domain and one Sterile Alpha Motif (SAM) domain. EPHB1 is able to stimulate fibroblast motility on extracellular matrix in a kinase-dependent manner, which is also correlated with its association with Grb7, an adaptor molecule implicated in the regulation of cell migration. It binds to Ephrin-B1, Ephrin-B2 and Ephrin-B3. EPHB1 plays an important roles in diverse biological processes including nervous system development, angiogenesis, and neural synapsis formation and maturation and may be involved in cell-cell interactions in the nervous system.

### Product Info

**Amount :** Fc) / 50 µg

**Content :** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.

**Storage condition :** Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.

**Amino Acid :** MEETLMDTRTATAELGWTANPASGWEEVSGYDENLNTIRTYQVCNVFEPNQNNWLLTTFINRRGA  
HRIYTEMRFTVRDCSSLPNVPGSCKETFNLYYYETDSVIATKKSAFWSEAPYLKVDITIAADESFSQV  
DFGGRLMKVNTEVRSFGPLTRNGFYLAQDYGACMSLLSVRVFFKKCPSIVQNFVFPETMTGAE  
STSLVIARGTCIPNAEEVDVPIKLYCNGDGEWMVPIGRCTCKPGYEPENSVACKACPAGTFKASQE  
AEGCSHCPSNSRSPAEASPICTCRTGYRADFPPEVACTSVPSGPRNVISIVNETSIILEWHPPRE  
TGGRDDVTYNIICKKCRADRRSCSRCDNVEFVPRQLGLETCRVSISLWAHTPYTFDIQAINGVSS  
KSPFPPQHVSVNITTNAAPSTVPIMHQVSATMRSITLSWPQPEQPNGIILDYEIRYYEKEHNEFNS  
SMARSQTNTARIDGLRPGMVVYVQVRARTVAGYGKFGSKMCFQTLTDDDYKSELREQLPVDDIEG  
RMDEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW  
YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQP  
REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYS  
KLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK

### Application Note

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 Åµg/ml. Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**Endotoxin :** Less than 0.1 ng/Åµg (1 IEU/Åµg) as determined by LAL test.